

## **SHORT COMMUNICATION**

# ROLE OF PLANT DOMINANTS ON ABANDONED TAILINGS CONTAINMENT FROM MANGANESE-ORE MAINING IN CHVALETICE, EASTERN BOHEMIA, CZECH REPUBLIC (OVERVIEW OF LONG-TERM CASE STUDIES)

MICHAL ŠTEFÁNEK

*Department of Biology, Faculty of Science, University of Hradec Králové, Rokitanského 62, 500 03 Hradec Králové*

*Department of Botany, Faculty of Science, Charles University in Prague, Benátská 2, 128 01 Praha 2; e-mail: stefim@seznam.cz*

**Received:** 9<sup>th</sup> November 2015, **Accepted:** 16<sup>th</sup> November 2015

## ABSTRACT

During long-term research (almost forty years) of tailings containment in Chvaletice (Eastern Bohemia) were also carried out studies on the role of each plant dominants in succession. This review presents the most interesting results of these studies.

**Keywords:** tailings containment, substrate toxicity, plant dominants, vegetation succession, colonization, lichens, bryophytes, annuals, grasses, pioneer tree species, species diversity

## INTRODUCTION

Tailings pond containment\* from manganese-ore maining in Chvaletice (Eastern Bohemia, Czech Republic) is a long-term target of studies on the succession of vegetation, but also the other biological processes and phenomena in extreme environmental conditions (high content of heavy metals and salts in the substrate, extreme fluctuations of surface temperatures, unfavourable water conditions). Studies of the systems of three sedimentation basins (working between 1952–1973) are underway with varying intensity, since the beginning of the phasing out the operation of the last tailings pond in 1973. Between 1986 and 2002 the research at this locality has been quite intensive (Kovář, 2004). First data concerning the local flora and vegetation after abandonment of the ore-waste deposits were published in the late 1970s (Kovář, 1977; 1979).

Two reclaimed tailings (1975–1978) demonstrate differences to the third one neighboring them. This unreclaimed abandoned waste body was left to spontaneous natural succession, that has been influenced only in places by anthropogenic activities (like deposition of construction wastes, motorcycling etc.). This enabled exposure to a number of experimental and monitoring studies, including those that were focused on the role of dominant plant species in primary succession, and contributed to some generalisations within the context of restoration ecology (e.g. Kovář, 1994; 1999; 2000; 2003; 2006; 2007; Prach & Hobbs, 2008; Prach *et al.*, 2008).

---

\* Terminological variations in literature could be found: abandoned (derelict) sedimentation basin, slime pit deposit, ore-washery setting pit, mining waste disposal, orewaste sulphate reservoir.

## MOSSES AND LICHENS

The initial vegetational cover of the substrate is characterised by the dominance of lichens of the genus *Cladonia*, mainly *Cladonia coniocraea* and *C. rei*, and also *Peltigera didactylus*. Besides lichens, the mosses that occur there include *Bryum caespiticium* and especially *Ceratodon purpureus* (Palice & Soldán, 2004; Soldán, 2009). After the fire episode (triggered by the extremely warm and dry summer in 1994), *Funaria hygrometrica* was increasingly abundant, but later its coverage decreased. Coverage of afore-mentioned species of lichens and mosses facilitate higher survival of clonal sprouts of dominant grasses (*Calamagrostis epigejos* and *Phragmites australis*), but it is unfavorable for long-term survival of seedlings. That is due to the microclimatic conditions (particularly soil humidity), which are suitable for seed germination, but not for further growth and attachment of seedlings. 75% of the seedlings of *Calamagrostis epigejos* in stands of mosses or lichens died within four months (Pohlová, 2004). Higher seedling establishment of this grass with prolonged success started after mulching of its own biomass (Štefánek *et al.*, 2012). As well as the terricolous species, some primary epiphytic lichens can sometimes grow on soil within the biological soil crust. In well-developed moss-lichen crust near the access road to the Chvaletice sedimentation basin, a prosperous thallus of *Evernia prunastri* was observed (Peksa, 2009). Much more preferable for the attachment of seedlings are crevices in stands of mosses, which extend deeper into the soil (see Hroudová & Zákavský, 2004). A similar role is also played by dynamic vegetation of mosses in dispersal facilitation of other plant dominants, which are able to disperse by seeds (f.e. *Betula pendula*). In Contrast, lichens are not suitable for attachment of seedlings (both genus *Cladonia* and species *Peltigera didactylus*). Rather surprisingly, the largest number of seedlings was observed on bare soil.

## ANNUAL PLANTS

Annual plant species (especially the vetches of the genus *Vicia* - *Vicia hirsuta* and *Vicia tetrasperma*) have appeared mainly in the younger stages of succession. Their proliferation and maximum abundance occurred within a relatively short period after the fire, which impacted a considerable area of the sedimentation basin in 1994. The sudden release of a high concentration of nutrients (and space) resulted in breaking the dormancy of seeds (Štefánek, 2004). A significant supporting factor of their dispersal was found in myrmecochory - dispersal of seeds by ants (Kovář *et al.*, 2013). The role of annual *Vicia* species was in enrichment of the soil with nitrogen through root-nodule bacteria.

## GRASSES

The main herb dominants in the middle successional stages are grasses - wood-small reed (*Calamagrostis epigejos*) within dry successional series and common reed (*Phragmites australis*) within wet successional series. Competitively strong and stress-tolerant *Calamagrostis epigejos* produces a positive effect in extreme environmental conditions, because it favorably affects seedling establishment of other plants due to its litter. The litter layer of *Calamagrostis epigejos* has a positive effect on the microclimate of the soil surface. Similarly, although to a lesser extent, common reed (*Calamagrostis epigejos*) may grow in wet places (in terrain depressions). Simultaneously, litter of *Calamagrostis epigejos* is, surprisingly, most easily subjected to decomposition (in comparison with other dominants – namely birch and aspen) (Štefánek *et al.*, 2012). Another comparative study of two types of

substrates deposited in sedimentation basins (ash deposits and ore-washery deposits) show noteworthy plasticity of this grass in its growth strategy (Bryndová & Kovář, 2004). It consists in different ways of dispersal and generative production of clones and their selection by the site conditions (Kovář *et al.*, 2004). Seed persistence in harsh environment on ecotoxicologically strong substrate is another supporting factor of vitality of *Calamagrostis epigeios* (Dostál & Kovář, 2013). The other grasses follow the stressed microsite conditions with their tolerance to extremes - e.g., *Puccinellia distans*, *Agrostis stolonifera*, *Festuca brevipila* etc. (Vaňková & Kovář, 2004).

## TREES

The dominant tree species are pioneer trees like birch (*Betula pendula*) and common aspen (*Populus tremula*), each of them with a different dispersal strategy. Birch spreads generative with small, anemochory seeds, while aspen is capable of clonal growth (but it is also possible to spread generatively with seed). Accordingly, they are also able to respond to different stresses and disturbances, such as sulphuric compounds (Rauch, 2004), or fire (Štefánek, 2004, Kovář *et al.*, 2011, Štefánek *et al.*, 2012). Effect upon leaf litter of these tree species is inconsiderable (with a similar effect as above-mentioned grasses), and it appears that leaf litter decay of both species occurs approximately at the same rate (Štefánek, 2004).

## CONCLUSIONS

Studies of different types of industrial waste in abandoned sedimentation deposits within the landscape, shows that extremely high concentrations of sulphuric compounds and heavy metals represent environmental conditions implying relatively low diversity of plants, and dominance of stress-tolerant and disturbance resistant plants (Vaňková & Kovář, 2004; Štefánek, 2004; Prach *et al.*, 2008). Described conditions are comparable with analogical environments throughout the world (such as volcanic substrates on high mountains) where some dominants are the same (e.g. bryophytes) regardless of latitude or longitude (Kovář, 2001). Anemochory and zoochory are the main types of plant diaspore distribution during plant colonisation of the substrate deposits, and they are the main factor of increasing biodiversity during spontaneous vegetational succession, when enrichment by humus and nutrients from decomposition processes of sufficient dead biomass is ensured (Kovář *et al.*, 2011; Štefánek *et al.*, 2012).

## ACKNOWLEDGEMENTS

The contribution was created with the institutional support of the Faculty of Science, University of Hradec Králové (project SV no. 2116) and Charles University in Prague. A big thanks to Pavel Kovář for valuable comments on the text.

## REFERENCES

Bryndová, I., & Kovář, P., (2004). Dynamics of the demographic parameters of the clonal plant *Calamagrostis epigeios* (L.) Roth in two kinds of industrial deposits (Abandoned sedimentation basins in Bukovina and Chvaletice). In: Kovář P. [ed.]: *Natural Recovery of Human-Made Deposits in Landscape* (pp. 267–276), (Biotic Interactions and Ore/Ash-Slag

Artificial Ecosystems), Academia, Prague.

Dostál, P., & Kovář, P., (2013). Seed rain and seed persistence of *Calamagrostis epigejos*. (L.) Roth in extreme ecotoxicological conditions as an abandoned ore-washery sedimentation basin. *Journal of Landscape Ecology*, 6(2): 17–33.

Hroudová, Z., Zákravský, P., (2004). The influence of the moss layer on soil surface microclimate in an abandoned ore-washery sedimentation basin. In: Kovář P. [ed.]: *Natural Recovery of Human-Made Deposits in Landscape* (pp. 235–247), (Biotic Interactions and Ore/Ash-Slag Artificial Ecosystems), Academia, Prague.

Kovář, P., (1977). Příspěvek ke květeně jihovýchodního Polabí. [Contribution to the flora of south-eastern Labe River Basin.]. *Zpr. Čs. Bot. Společ.*, 12: 83–92. [in Czech]

Kovář, P., (1979). Geobotanické aspekty rekultivace odkališť manganorudných a kyzových závodů Chvaletice. Ecological aspects of slime pit recultivation of the MKK-Chvaletice. *Práce a Studie-Prír.*, 11: 63–78. [in Czech]

Kovář, P., (1981). Společenstvo s *Conium maculatum* na odkalištích MKZ Chvaletice. [Plant community of *Conium maculatum* on the sedimentation pond deposits at Chvaletice.] – *Zpr. Čs. Bot. Společ.*, 15: 111–112. [in Czech]

Kovář, P., (1994). Vegetační monitorování a ekologie obnovy krajiny: změny na odkalištích MKZ Chvaletice. [Vegetation monitoring and restoration ecology in landscape: changes of sediment deposits at Chvaletice.] – In: Kirschnerová, L., [ed.]: *Monitoring vybraných přirozených společenstev a populací rostlinných indikátorů v České republice*. [Monitoring in selected natural communities and populations of plant indicators in the Czech Republic.] – *Příroda*, Praha, 1: 79–96. [in Czech]

Kovář, P., (1999). Biotic interactions and restoration ecology of abandoned sedimentation ponds of toxic materials. – In: Kovář P. [ed.]: *Nature and Culture in Landscape Ecology* (pp. 290–302), (Experiences for the 3rd Millennium), The Karolinum Press, Praha.

Kovář, P., (2003). Transplantovaná synantropizace: rostliny a nová stanoviště vložená do krajiny. [Synanthropisation transplanted: plants and new habitats introduced into landscape.] – *Zpr. Čs. Bot. Společ.* 38, Mater. 19: 17–26. [in Czech]

Kovář, P., [ed.] (2004). *Natural Recovery of Human-Made Deposits in Landscape* (Biotic Interactions and Ore/Ash-Slag Artificial Ecosystems), Academia, Prague, 2004, 358 p.

Kovář, P., (2006). Ekologie obnovy poškozené krajiny. [Restoration ecology of damaged landscape.] – *Zpr. Čs. Bot. Společ.*, Praha, 41, Mater. 21: 23–38. [in Czech]

Kovář, P., (2000). Abandoned anthropogenic landscapes: Are they potentially multifunctional? – In: Brandt J., Tress B. et Tress G. [eds]: *Multifunctional landscapes. Interdisciplinary approaches to landscape research and management* (p. 251). Conference Material, Centre for Landscape Res., Roskilde, Denmark.

Kovář, P., (2001). Young landscapes: Ecological phenomena on South-American volcanoes and Central-European industrial substrates. – In: Mander, Ú., Printsman, A. et Palang, H., [eds]: *Conf. Proc. "Development of European Landscapes"* (pp. 529–231), IALE European Conf. 2001, Vol. II., Publicationes Inst. Geogr., Universitatis Tartuensis, 92.

Kovář, P., (2007). Terrestrial islands of non-reclaimed industrial deposits: plant species diversity and cultural landscape. – In: Bunce, R.G.H., Jongman, R.H.G., Hojas, L., Weei, S.: *25 years of landscape ecology: scientific principles in practice* (pp. 694–695). Proc. of the 7th IALE World Congress – Part 2, Wageningen, The Netherlands, 8–12 July, 2007, Theme 6: Cultural landscapes and landscape management. 6.2 Symposium 20: Landscape diversity and biological diversity cultural landscapes.

- Kovář, P., Štěpánek, J. et Kirschner, J., (2004). Clonal diversity of *Calamagrostis epigejos* (L.) Roth in relation to type of industrial substrate and successional stage. – In: Kovář, P., [ed.]: *Natural Recovery of Human-Made Deposits in Landscape* (pp. 285–293), (Biotic Interactions and Ore/Ash-Slag Artificial Ecosystems), Academia, Prague.
- Kovář, P., Štěfánek, M. et Mrázek, J., (2011). Responses of vegetation stages with woody dominants to stress and disturbance during succession on abandoned tailings in cultural landscape. *Journal of Landscape Ecology*, 4/2: 35–48.
- Kovář, P., Vojtíšek, P. et Zentsová, I., (2013). Ants as ecosystem engineers in natural restoration of human made habitats. *Journal of Landscape Ecology*, 6(1): 18–31.
- Palice, Z. & Soldán, Z., (2004). Lichen and bryophyte species diversity on toxic substrates in the abandoned sedimentation basins of Chvaletice and Bukovina. – In: Kovář, P., [ed.]: *Natural Recovery of Human-Made Deposits in Landscape* (pp. 200–221), (Biotic Interactions and Ore/Ash-Slag Artificial Ecosystems). Academia, Prague.
- Peksa, O., (2009). 3.5. Lichens – species composition and diversity. – In: Neustupa, J., Škaloud, P., Peksa, O., Kubátová, A., Soldán, Z., Černá, K., Prášil, K., Bukovská, P., Vojta, J., Pažoutová, M., Veselá, J., Škaloudová, M.: *The biological soil crusts in Central European ecosystems, with special reference to taxonomic structure and ecology of the surface crusts at Czech ore-waste and ash-slag sedimentation industrial basins* (pp. 19–21) – Novit. Bot. Univ. Carol. 19/2008.
- Pohlová, R., (2004). Changes on microsites of the moss *Ceratodon purpureus* and lichens *Peltigera didactyla* and *Cladonia* sp. div. in the abandoned sedimentation basin in Chvaletice. – In: Kovář, P., [ed.]: *Natural Recovery of Human-Made Deposits in Landscape* (pp. 222–234), (Biotic Interactions and Ore/Ash-Slag Artificial Ecosystems). Academia, Prague.
- Prach, K., Bastl, M., Konvalinková, P., Kovář, P., Novák, J., Pyšek, P., Řehouňková, K. et Sádlo, J., (2008). Sukcese vegetace na antropogenních stanovištích v České republice – přehled dominantních druhů a stadií. [Vegetation succession in human-made habitats in the Czech Republic – survey of dominant species and stages.] – *Příroda*, 26: 5–26. Praha. [in Czech]
- Prach, K. & Hobbs, R. J., (2008). Spontaneous succession versus technical reclamation in the restoration of disturbed sites. *Restoration Ecology* 16(3): 363–366.
- Soldán, Z., (2009). 3.7. Species composition and diversity of bryophytes on anthropogenic substrata. – In: Neustupa, J., Škaloud, P., Peksa, O., Kubátová, A., Soldán, Z., Černá, K., Prášil, K., Bukovská, P., Vojta, J., Pažoutová, M., Veselá, J. et Škaloudová, M.: *The biological soil crusts in Central European ecosystems, with special reference to taxonomic structure and ecology of the surface crusts at Czech ore-waste and ash-slag sedimentation industrial basins* (pp. 45–50).– Novit. Bot. Univ. Carol. 19/2008.
- Rauch, O., (2004): Genesis and characteristics of orewaste sulphate soils at Chvaletice. – In: Kovář, P., [ed.]: *Natural Recovery of Human-Made Deposits in Landscape* (pp. 45–58), (Biotic Interactions and Ore/Ash-Slag Artificial Ecosystems), Academia, Prague.
- Štěfánek, M., (2004): Secondary succession after fire on an abandoned ore-washery sedimentation basin – different trajectories (A comparison with primary succession). – In: Kovář, P., [ed.]: *Natural Recovery of Human-Made Deposits in Landscape* (pp. 248–266 ), (Biotic Interactions and Ore/Ash-Slag Artificial Ecosystems), Academia, Prague.

Štefánek, M., Kovář, P., & Dlouhá, V., (2012): Role of fire episode, leaf litter decomposition and mulching effects in restoration of surface soil crust microecosystem on abandoned tailings containment. *Journal of Landscape Ecology*, 5/3: 57–69.

Vaňková, J., & Kovář, P., (2004): Plant species diversity in the biotopes of un-reclaimed industrial deposits as artificial islands in landscape. – In: Kovář, P., [ed.]: *Natural Recovery of Human-Made Deposits in Landscape* (pp. 30–45), (Biotic Interactions and Ore/Ash-Slag Artificial Ecosystems). Academia, Prague.